

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Ethoxy-6-([2-[(3-ethoxy-2-hydroxy-benzylidene)amino]benzyl]iminomethyl)phenol

K. U. Ambili,^a S. S. Sreejith,^a Jinsa Mary Jacob,^a M. Sithambaresan^{b*} and M. R. Prathapachandra Kurup^a^aDepartment of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, India, and ^bDepartment of Chemistry, Faculty of Science, Eastern University, Sri Lanka, Chenkalady, Sri Lanka
Correspondence e-mail: eesans@yahoo.com


Received 27 June 2012; accepted 10 July 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}—\text{C}) = 0.003$ Å; R factor = 0.038; wR factor = 0.112; data-to-parameter ratio = 13.2.

The title compound, $\text{C}_{25}\text{H}_{26}\text{N}_2\text{O}_4$, exists in an *E* conformation with respect to each azomethine link. The two phenol-substituted benzene rings are twisted away from the plane of the diimine benzene ring by dihedral angles of 27.25 (5) and 56.67 (5)°. The molecular structure is stabilized by intra-molecular $\text{O}—\text{H} \cdots \text{N}$ hydrogen bonds.

Related literature

For the applications of salen Schiff bases, see: Cozzi (2004); Hodnett & Dunn (1970). For the synthesis of Schiff bases, see: Tümer (2000). For a related structure, see: Aslantaş *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{25}\text{H}_{26}\text{N}_2\text{O}_4$
 $M_r = 418.48$
Monoclinic, $P2_1/c$
 $a = 4.8315$ (4) Å $b = 17.5414$ (14) Å
 $c = 25.828$ (2) Å
 $\beta = 94.356$ (3)°
 $V = 2182.6$ (3) Å³ $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹ $T = 296$ K
 $0.40 \times 0.20 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 2004)
 $T_{\min} = 0.979$, $T_{\max} = 0.983$ 32150 measured reflections
3833 independent reflections
2543 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.112$
 $S = 1.05$
3833 reflections
291 parameters
2 restraintsH atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.13$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{O2}—\text{H2O} \cdots \text{N1}$	0.85 (1)	1.79 (1)	2.5711 (19)	151 (2)
$\text{O3}—\text{H3O} \cdots \text{N2}$	0.85 (1)	1.80 (1)	2.5844 (19)	152 (2)

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97 and publCIF (Westrip, 2010).

The authors are grateful to the Sophisticated Analytical Instruments Facility, Cochin University of Science and Technology, Kochi-22, India, for providing the single-crystal X-ray diffraction data. MRPK, KUA and SSS thank the Defence Research Development Organization, New Delhi, India, for financial support. JMJ thanks the Council of Scientific and Industrial Research, New Delhi, India, for a Senior Research Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2580).

References

- Aslantaş, M., Tümer, M., Şahin, E. & Tümer, F. (2007). *Acta Cryst.* **E63**, o644–o645.
- Bruker (2004). *SADABS*, *APEX2*, *XPREP* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cozzi, P. G. (2004). *Chem. Soc. Rev.* **33**, 410–421.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hodnett, E. M. & Dunn, W. J. (1970). *J. Med. Chem.* **13**, 768–770.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tümer, M. (2000). *Synth. React. Inorg. Met. Org. Chem.* **30**, 1139–1158.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o2482 [doi:10.1107/S1600536812031479]

2-Ethoxy-6-({2-[(3-ethoxy-2-hydroxybenzylidene)amino]benzyl}iminomethyl)-phenol

K. U. Ambili, S. S. Sreejith, Jinsa Mary Jacob, M. Sithambaresan and M. R. Prathapachandra Kurup

Comment

The chelating structure, moderate electron donation and easy tunable electronic and steric effects make dicompartamental salen type Schiff bases to act as versatile ligands. They are able to stabilize different metals in various oxidation states and control the performance of the metals in various catalytic transformations (Cozzi, 2004). Schiff bases are very selective in the sense that they provide geometrical cavity control for host guest interaction and modulate its lipophilicity to stabilize a specific metal ion. Moreover, it has been suggested that the azomethine linkage in Schiff bases is responsible for their biological activities such as antimicrobial, antifungal and antitumor and also to be used as herbicides (Hodnett & Dunn, 1970).

The *E* conformation of the compound is evidenced from the torsion angles, 176.94 (14)° and 179.38 (15)° made by the C10—N1—C9—C7 and C16—N2—C17—C18 linkages respectively. The bond lengths and bond angles are in normal ranges and agree with the related structure (Aslantaş, *et al.*, 2007). The crystal involves two intramolecular O—H⋯N hydrogen bonds and C—H⋯ π interactions which make the molecule stable.

Experimental

The title compound was prepared according to the reported procedure (Tümer, 2000) by the condensation of the ethanolic solution of 3-ethoxy-2-hydroxybenzaldehyde (1 mmol, 0.166 g) with an ethanolic solution of 2-aminobenzylamine (0.5 mmol, 0.061 g). The reaction mixture was heated to reflux for 6 h and kept for cooling at room temperature. The slow evaporation yielded orange-yellow crystals of *N,N'*-bis(3-ethoxy-2-hydroxybenzylidene)-2-aminobenzylamine.

Refinement

All H atoms on C were placed in calculated positions, guided by difference maps, with C—H bond distances 0.93–0.97 Å. H atoms were assigned as $U_{\text{iso}}=1.2U_{\text{eq}}$ (1.5 for Me). O(2)—H(2o) and O(3)—H(3o) H atoms were located from difference maps and restrained using *DFIX* instructions.

Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREF* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *pubCIF* (Westrip, 2010).




Figure 1

ORTEP view of the unique part of the compound, drawn with 50% probability displacement ellipsoids for the non-H atoms.




Figure 2

Packing diagram of the compound viewed along *a* axis.

2-Ethoxy-6-({2-[(3-ethoxy-2-hydroxybenzylidene)amino]benzyl}iminomethyl)phenol

Crystal data

$C_{25}H_{26}N_2O_4$	$F(000) = 888$
$M_r = 418.48$	$D_x = 1.273 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2_1bc$	Cell parameters from 5446 reflections
$a = 4.8315 (4) \text{ \AA}$	$\theta = 2.3\text{--}22.4^\circ$
$b = 17.5414 (14) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 25.828 (2) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 94.356 (3)^\circ$	Needle-like, orange
$V = 2182.6 (3) \text{ \AA}^3$	$0.40 \times 0.20 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII CCD	32150 measured reflections
diffractometer	3833 independent reflections
Radiation source: fine-focus sealed tube	2543 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.044$
ω and ϕ scan	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan	$h = -5 \rightarrow 5$
(<i>SADABS</i> ; Bruker, 2004)	$k = -20 \rightarrow 20$
$T_{\text{min}} = 0.979$, $T_{\text{max}} = 0.983$	$l = -30 \rightarrow 30$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.038$	$w = 1/[\sigma^2(F_o^2) + (0.0493P)^2 + 0.3451P]$
$wR(F^2) = 0.112$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3833 reflections	$\Delta\rho_{\text{max}} = 0.13 \text{ e \AA}^{-3}$
291 parameters	$\Delta\rho_{\text{min}} = -0.13 \text{ e \AA}^{-3}$
2 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0062 (9)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.5523 (3)	1.08404 (8)	0.43051 (5)	0.0645 (4)
O2	1.1570 (3)	1.09379 (8)	0.35604 (5)	0.0598 (4)
O3	0.0532 (3)	0.93676 (8)	0.17006 (5)	0.0543 (4)

O4	−0.3152 (3)	0.82995 (8)	0.15297 (5)	0.0659 (4)
N1	0.8660 (3)	1.03221 (9)	0.27970 (5)	0.0466 (4)
N2	0.4050 (3)	1.03122 (8)	0.13518 (5)	0.0445 (4)
C1	1.8136 (6)	1.16138 (15)	0.48999 (10)	0.0968 (9)
H1A	1.8577	1.1960	0.4630	0.145*
H1B	1.9602	1.1620	0.5172	0.145*
H1C	1.6429	1.1768	0.5037	0.145*
C2	1.7824 (4)	1.08294 (13)	0.46832 (8)	0.0634 (6)
H2A	1.9497	1.0679	0.4524	0.076*
H2B	1.7496	1.0469	0.4957	0.076*
C3	1.4950 (4)	1.01888 (11)	0.40288 (7)	0.0479 (5)
C4	1.6259 (4)	0.94994 (12)	0.41106 (8)	0.0589 (5)
H4	1.7682	0.9453	0.4372	0.071*
C5	1.5485 (5)	0.88751 (12)	0.38081 (8)	0.0669 (6)
H5	1.6380	0.8411	0.3869	0.080*
C6	1.3413 (4)	0.89335 (12)	0.34193 (8)	0.0582 (5)
H6	1.2913	0.8510	0.3217	0.070*
C7	1.2044 (3)	0.96269 (10)	0.33256 (6)	0.0433 (4)
C8	1.2814 (3)	1.02555 (10)	0.36311 (7)	0.0431 (4)
C9	0.9881 (4)	0.96941 (11)	0.29046 (7)	0.0464 (5)
H9	0.9376	0.9265	0.2708	0.056*
C10	0.6616 (3)	1.03497 (11)	0.23529 (6)	0.0471 (5)
H10A	0.4828	1.0494	0.2470	0.057*
H10B	0.6430	0.9847	0.2198	0.057*
C11	0.7426 (3)	1.09114 (10)	0.19477 (7)	0.0413 (4)
C12	0.9465 (4)	1.14566 (11)	0.20497 (7)	0.0503 (5)
H12	1.0399	1.1475	0.2378	0.060*
C13	1.0149 (4)	1.19732 (11)	0.16787 (8)	0.0583 (5)
H13	1.1529	1.2334	0.1756	0.070*
C14	0.8783 (4)	1.19519 (11)	0.11950 (8)	0.0610 (6)
H14	0.9212	1.2305	0.0945	0.073*
C15	0.6775 (4)	1.14086 (11)	0.10774 (7)	0.0550 (5)
H15	0.5866	1.1394	0.0747	0.066*
C16	0.6101 (3)	1.08823 (10)	0.14499 (7)	0.0419 (4)
C17	0.3525 (4)	1.00201 (11)	0.09022 (7)	0.0468 (5)
H17	0.4510	1.0191	0.0629	0.056*
C18	0.1456 (3)	0.94344 (10)	0.08038 (7)	0.0438 (4)
C19	0.0901 (4)	0.91471 (12)	0.03008 (7)	0.0565 (5)
H19	0.1824	0.9347	0.0028	0.068*
C20	−0.0987 (4)	0.85754 (13)	0.02086 (8)	0.0636 (6)
H20	−0.1334	0.8385	−0.0126	0.076*
C21	−0.2384 (4)	0.82791 (12)	0.06090 (9)	0.0604 (6)
H21	−0.3671	0.7891	0.0541	0.072*
C22	−0.1905 (4)	0.85472 (11)	0.11067 (7)	0.0493 (5)
C23	0.0053 (3)	0.91322 (10)	0.12070 (7)	0.0430 (4)
C24	−0.4979 (4)	0.76620 (12)	0.14644 (10)	0.0709 (6)
H24A	−0.3996	0.7224	0.1341	0.085*
H24B	−0.6513	0.7780	0.1213	0.085*
C25	−0.6032 (6)	0.74931 (17)	0.19800 (12)	0.1079 (10)

H25A	−0.4492	0.7407	0.2230	0.162*
H25B	−0.7181	0.7046	0.1954	0.162*
H25C	−0.7102	0.7918	0.2088	0.162*
H2O	1.031 (4)	1.0877 (14)	0.3316 (6)	0.090 (8)*
H3O	0.172 (4)	0.9724 (10)	0.1691 (10)	0.103 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0646 (9)	0.0605 (9)	0.0640 (9)	0.0055 (7)	−0.0227 (7)	−0.0080 (7)
O2	0.0650 (9)	0.0476 (9)	0.0632 (9)	0.0063 (7)	−0.0189 (7)	−0.0030 (7)
O3	0.0597 (8)	0.0607 (9)	0.0431 (8)	−0.0100 (7)	0.0075 (6)	−0.0064 (7)
O4	0.0606 (8)	0.0640 (10)	0.0739 (10)	−0.0166 (7)	0.0109 (7)	−0.0022 (8)
N1	0.0454 (8)	0.0537 (10)	0.0400 (8)	−0.0027 (7)	−0.0006 (7)	0.0008 (7)
N2	0.0416 (8)	0.0480 (9)	0.0432 (9)	0.0004 (7)	−0.0005 (6)	0.0010 (7)
C1	0.115 (2)	0.0734 (18)	0.0933 (19)	−0.0083 (15)	−0.0501 (16)	−0.0043 (15)
C2	0.0626 (12)	0.0727 (16)	0.0519 (12)	−0.0012 (11)	−0.0152 (10)	0.0006 (11)
C3	0.0471 (10)	0.0514 (13)	0.0448 (11)	−0.0009 (9)	−0.0006 (8)	0.0009 (9)
C4	0.0594 (12)	0.0619 (14)	0.0536 (12)	0.0102 (10)	−0.0080 (10)	0.0044 (11)
C5	0.0766 (14)	0.0535 (14)	0.0691 (14)	0.0177 (11)	−0.0043 (12)	0.0057 (12)
C6	0.0705 (13)	0.0456 (12)	0.0576 (13)	0.0023 (10)	0.0004 (10)	−0.0035 (10)
C7	0.0465 (10)	0.0436 (11)	0.0400 (10)	−0.0036 (8)	0.0043 (8)	0.0056 (9)
C8	0.0451 (10)	0.0412 (11)	0.0428 (10)	0.0020 (8)	0.0031 (8)	0.0063 (9)
C9	0.0510 (10)	0.0495 (12)	0.0391 (10)	−0.0096 (9)	0.0060 (8)	−0.0006 (9)
C10	0.0421 (9)	0.0580 (12)	0.0405 (10)	−0.0053 (8)	−0.0009 (8)	0.0031 (9)
C11	0.0383 (9)	0.0430 (10)	0.0425 (10)	0.0024 (8)	0.0036 (7)	0.0001 (8)
C12	0.0482 (10)	0.0503 (12)	0.0517 (11)	−0.0047 (9)	0.0001 (9)	−0.0025 (10)
C13	0.0593 (12)	0.0485 (12)	0.0672 (14)	−0.0123 (10)	0.0059 (10)	−0.0003 (10)
C14	0.0736 (14)	0.0491 (13)	0.0609 (14)	−0.0082 (10)	0.0093 (11)	0.0105 (10)
C15	0.0641 (12)	0.0540 (13)	0.0463 (11)	−0.0020 (10)	−0.0005 (9)	0.0061 (10)
C16	0.0388 (9)	0.0429 (11)	0.0440 (10)	0.0007 (8)	0.0030 (8)	0.0002 (8)
C17	0.0482 (10)	0.0528 (12)	0.0394 (11)	0.0020 (9)	0.0032 (8)	0.0031 (9)
C18	0.0439 (9)	0.0472 (11)	0.0394 (10)	0.0044 (8)	−0.0025 (8)	−0.0033 (9)
C19	0.0628 (12)	0.0649 (14)	0.0411 (11)	0.0010 (11)	−0.0010 (9)	−0.0038 (10)
C20	0.0668 (13)	0.0709 (15)	0.0508 (13)	0.0024 (12)	−0.0115 (10)	−0.0182 (11)
C21	0.0501 (11)	0.0577 (13)	0.0712 (15)	−0.0023 (10)	−0.0094 (10)	−0.0147 (12)
C22	0.0408 (9)	0.0494 (12)	0.0573 (13)	0.0023 (9)	0.0011 (9)	−0.0023 (10)
C23	0.0402 (9)	0.0468 (11)	0.0414 (11)	0.0059 (8)	−0.0009 (8)	−0.0045 (9)
C24	0.0572 (12)	0.0480 (13)	0.1072 (19)	−0.0059 (10)	0.0038 (12)	0.0088 (12)
C25	0.099 (2)	0.093 (2)	0.134 (2)	−0.0265 (16)	0.0211 (18)	0.0369 (18)

Geometric parameters (\AA , $^\circ$)

O1—C3	1.365 (2)	C10—H10A	0.9700
O1—C2	1.423 (2)	C10—H10B	0.9700
O2—C8	1.346 (2)	C11—C12	1.384 (2)
O2—H2O	0.8502 (11)	C11—C16	1.393 (2)
O3—C23	1.343 (2)	C12—C13	1.377 (3)
O3—H3O	0.8501 (11)	C12—H12	0.9300
O4—C22	1.358 (2)	C13—C14	1.368 (3)

O4—C24	1.427 (2)	C13—H13	0.9300
N1—C9	1.270 (2)	C14—C15	1.377 (3)
N1—C10	1.456 (2)	C14—H14	0.9300
N2—C17	1.277 (2)	C15—C16	1.390 (2)
N2—C16	1.417 (2)	C15—H15	0.9300
C1—C2	1.489 (3)	C17—C18	1.443 (2)
C1—H1A	0.9600	C17—H17	0.9300
C1—H1B	0.9600	C18—C23	1.390 (2)
C1—H1C	0.9600	C18—C19	1.400 (2)
C2—H2A	0.9700	C19—C20	1.364 (3)
C2—H2B	0.9700	C19—H19	0.9300
C3—C4	1.374 (3)	C20—C21	1.379 (3)
C3—C8	1.404 (2)	C20—H20	0.9300
C4—C5	1.380 (3)	C21—C22	1.372 (3)
C4—H4	0.9300	C21—H21	0.9300
C5—C6	1.367 (3)	C22—C23	1.407 (2)
C5—H5	0.9300	C24—C25	1.491 (4)
C6—C7	1.397 (3)	C24—H24A	0.9700
C6—H6	0.9300	C24—H24B	0.9700
C7—C8	1.390 (2)	C25—H25A	0.9600
C7—C9	1.455 (2)	C25—H25B	0.9600
C9—H9	0.9300	C25—H25C	0.9600
C10—C11	1.510 (2)		
C3—O1—C2	117.69 (15)	C13—C12—C11	121.82 (18)
C8—O2—H2O	106.0 (17)	C13—C12—H12	119.1
C23—O3—H3O	105.4 (18)	C11—C12—H12	119.1
C22—O4—C24	117.63 (16)	C14—C13—C12	119.52 (18)
C9—N1—C10	118.48 (16)	C14—C13—H13	120.2
C17—N2—C16	122.26 (15)	C12—C13—H13	120.2
C2—C1—H1A	109.5	C13—C14—C15	120.20 (19)
C2—C1—H1B	109.5	C13—C14—H14	119.9
H1A—C1—H1B	109.5	C15—C14—H14	119.9
C2—C1—H1C	109.5	C14—C15—C16	120.34 (18)
H1A—C1—H1C	109.5	C14—C15—H15	119.8
H1B—C1—H1C	109.5	C16—C15—H15	119.8
O1—C2—C1	107.27 (18)	C15—C16—C11	119.93 (16)
O1—C2—H2A	110.3	C15—C16—N2	122.85 (16)
C1—C2—H2A	110.3	C11—C16—N2	117.21 (15)
O1—C2—H2B	110.3	N2—C17—C18	122.22 (16)
C1—C2—H2B	110.3	N2—C17—H17	118.9
H2A—C2—H2B	108.5	C18—C17—H17	118.9
O1—C3—C4	125.73 (17)	C23—C18—C19	119.22 (17)
O1—C3—C8	114.93 (16)	C23—C18—C17	120.68 (16)
C4—C3—C8	119.35 (18)	C19—C18—C17	120.07 (17)
C3—C4—C5	120.62 (18)	C20—C19—C18	120.32 (19)
C3—C4—H4	119.7	C20—C19—H19	119.8
C5—C4—H4	119.7	C18—C19—H19	119.8
C6—C5—C4	120.56 (19)	C19—C20—C21	120.34 (19)

C6—C5—H5	119.7	C19—C20—H20	119.8
C4—C5—H5	119.7	C21—C20—H20	119.8
C5—C6—C7	120.20 (19)	C22—C21—C20	121.08 (19)
C5—C6—H6	119.9	C22—C21—H21	119.5
C7—C6—H6	119.9	C20—C21—H21	119.5
C8—C7—C6	119.32 (16)	O4—C22—C21	126.19 (18)
C8—C7—C9	120.42 (16)	O4—C22—C23	114.74 (16)
C6—C7—C9	120.25 (17)	C21—C22—C23	119.07 (18)
O2—C8—C7	122.08 (15)	O3—C23—C18	122.37 (16)
O2—C8—C3	117.98 (16)	O3—C23—C22	117.65 (16)
C7—C8—C3	119.95 (17)	C18—C23—C22	119.97 (16)
N1—C9—C7	121.94 (17)	O4—C24—C25	107.5 (2)
N1—C9—H9	119.0	O4—C24—H24A	110.2
C7—C9—H9	119.0	C25—C24—H24A	110.2
N1—C10—C11	111.81 (14)	O4—C24—H24B	110.2
N1—C10—H10A	109.3	C25—C24—H24B	110.2
C11—C10—H10A	109.3	H24A—C24—H24B	108.5
N1—C10—H10B	109.3	C24—C25—H25A	109.5
C11—C10—H10B	109.3	C24—C25—H25B	109.5
H10A—C10—H10B	107.9	H25A—C25—H25B	109.5
C12—C11—C16	118.15 (16)	C24—C25—H25C	109.5
C12—C11—C10	122.49 (16)	H25A—C25—H25C	109.5
C16—C11—C10	119.36 (15)	H25B—C25—H25C	109.5
C3—O1—C2—C1	175.91 (19)	C14—C15—C16—C11	1.1 (3)
C2—O1—C3—C4	5.6 (3)	C14—C15—C16—N2	179.84 (17)
C2—O1—C3—C8	−174.61 (16)	C12—C11—C16—C15	−2.1 (2)
O1—C3—C4—C5	179.47 (19)	C10—C11—C16—C15	178.31 (17)
C8—C3—C4—C5	−0.4 (3)	C12—C11—C16—N2	179.06 (15)
C3—C4—C5—C6	0.5 (3)	C10—C11—C16—N2	−0.5 (2)
C4—C5—C6—C7	−0.3 (3)	C17—N2—C16—C15	30.6 (3)
C5—C6—C7—C8	0.1 (3)	C17—N2—C16—C11	−150.65 (17)
C5—C6—C7—C9	178.64 (18)	C16—N2—C17—C18	179.38 (15)
C6—C7—C8—O2	−179.86 (16)	N2—C17—C18—C23	−3.8 (3)
C9—C7—C8—O2	1.6 (3)	N2—C17—C18—C19	178.20 (17)
C6—C7—C8—C3	0.0 (3)	C23—C18—C19—C20	−0.2 (3)
C9—C7—C8—C3	−178.53 (15)	C17—C18—C19—C20	177.81 (17)
O1—C3—C8—O2	0.2 (2)	C18—C19—C20—C21	0.5 (3)
C4—C3—C8—O2	179.99 (17)	C19—C20—C21—C22	−0.3 (3)
O1—C3—C8—C7	−179.72 (15)	C24—O4—C22—C21	5.1 (3)
C4—C3—C8—C7	0.1 (3)	C24—O4—C22—C23	−174.61 (16)
C10—N1—C9—C7	176.94 (14)	C20—C21—C22—O4	−179.98 (19)
C8—C7—C9—N1	1.0 (3)	C20—C21—C22—C23	−0.2 (3)
C6—C7—C9—N1	−177.52 (17)	C19—C18—C23—O3	178.33 (16)
C9—N1—C10—C11	−119.33 (18)	C17—C18—C23—O3	0.3 (3)
N1—C10—C11—C12	−14.9 (2)	C19—C18—C23—C22	−0.3 (3)
N1—C10—C11—C16	164.62 (15)	C17—C18—C23—C22	−178.31 (16)
C16—C11—C12—C13	1.6 (3)	O4—C22—C23—O3	1.6 (2)
C10—C11—C12—C13	−178.91 (17)	C21—C22—C23—O3	−178.17 (16)

C11—C12—C13—C14	0.1 (3)	O4—C22—C23—C18	−179.72 (16)
C12—C13—C14—C15	−1.2 (3)	C21—C22—C23—C18	0.5 (3)
C13—C14—C15—C16	0.6 (3)	C22—O4—C24—C25	178.99 (18)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2O \cdots N1	0.85 (1)	1.79 (1)	2.5711 (19)	151 (2)
O3—H3O \cdots N2	0.85 (1)	1.80 (1)	2.5844 (19)	152 (2)